# Deriving Growing Season Metrics from MODIS EVI and a Simple Bioclimatic Index


JOANNE NIGHTINGALE<sup>2,8</sup>, NICHOLAS COOPS AND RICHARD WARING<sup>2</sup>

# OBJECTIVE


Test whether general phenological behaviour can be accurately characterised at monthly intervals using MODIS EVI data

### GROWING SEASON INDEX

- ♦ Index to quantify greenness of vegetation during the year [1] (Fig. 1)
- Uses 3 dimensionless climatic modifiers to account for variations observed in seasonal phenology (Fig. 2a-c)




# HOW ACCURATE IS THE GSI?



#### COMPARISON WITH MODIS EVI

- MODIS EVI satellite measure of vegetation greenness (Fig 4.)
- Analysis based on EPA Level II ecological zones across the USA and dominant MODIS derived IGBP class (forest / non-forest) (Fig. 3)
- Significant associations in seasonal phenology for forest as opposed to non-forest classes (Fig. 5)






Fig. 5: Seasonal greenness of forest versus non-forest land cover types depicted with the GSI and EVI





Fig. 1: Climatological limits to plant growth defined by the Growing Season Index[1]






Fig. 7: EVI-derived start of growing season for forests across the United States

# GROWING SEASON METRICS

 Thresholds to define Start and End of GSI and EVI growing seasons (Table 2 and Fig. 6)




Table 2: Thresholds applied to the GSI and EVI long-term average data to derive growing Season metrics

Threshold	Value	Description		
GSI	0.5 [1]	>0.5, sustained canopy		
EVI	Seasonal Mid-point [2]	Half-max of long-term av.		

- Fig. 6: Schemtic diagram showing derivation of growing season metrics using defined thresholds
- Poor association between GSI and independent measure (SM) of EVI growing season start, end and length (Table 3)

Table 3: Association between GSI and independent estimate of EVI growing season metrics

Veg Type	GSI Start	EVI Start	GSI End	EVI End	GSI Length	EVI Length
Forest	March	April	September	October	6 months	7 months
Non-Forest	January	February	September	November	7 months	4 months
R <sup>2</sup>	0.34		0.00		0.24	

- GSI used to provide thresholds to define EVI growing season attributes for forest (0.28) & non-forest (0.22) classes (Fig. 6)
- ♦ Improved relationship between GSI & EVI Start ( $R^2 = 1$ ), End ( $R^2 = 0.62$ ) and Length ( $R^2 = 0.35$ ) of forest growing season metrics (Fig. 7)


#### CONCLUSIONS

- Monthly phenological behaviour of forests can be accurately characterised using the MODIS EVI across the continental USA
- GSI and EVI are less useful in defining the growing season attributes of vegetation with low canopy densities (i.e. grassland and savannas)

# ACKNOWLEDGEMENTS & REFERENCES

We acknowledge support for this research from NASA (NNG04GK26G)

Nightingale, J. Coops, N. and Waring, R. Using a Monthly Bioclimatic Index with MODIS Vegetation Attributes to Estimate Seasonal Phenology across the Contiguous United States. Global Change Biology. (In press).



UBC

[1] Jolly, M. *et al.* 2005. A generalised, bioclimatic index to predict foliar phenology in response to climate. Global Change Biology. 11: 619-632.

[2] Schwartz, M.D. *et al.* 2002. Assessing satellite-derived start of season measures in the conterminous USA. International Journal of Climatology. 22: 1793-1805.





<sup>A</sup> College of Forestry, Oregon State University, Corvallis, USA

<sup>B</sup> Integrated Remote Sensing Studio, Forest Sciences Centre, University of British Columbia, Vancouver, Canada

Contact Author: Joanne.Nightingale@ubc.ca

